

Partial Supersymmetry Breaking from Five Dimensions

Richard Altendorfer

*Department of Physics and Astronomy
The Johns Hopkins University
3400 North Charles Street
Baltimore, MD 21218, U.S.A.*

Abstract

Theories of partial supersymmetry breaking $N = 2 \rightarrow N = 1$ in four dimensions are derived by coupling the $N = 2$ massless gravitino multiplet to $N = 2$ supergravity in five dimensions and performing a generalized dimensional reduction on S^1/Z_2 with the Scherk-Schwarz mechanism. These theories agree with results that were previously derived from four dimensions.

1 Introduction

Most supersymmetry phenomenology is based on the minimal supersymmetric standard model, which is assumed to be the low energy limit of a more fundamental theory like string theory or M-theory. Those fundamental theories are formulated in higher space-time dimensions and possess an extended number of supersymmetries which have to be spontaneously broken to arrive at a phenomenologically viable $N = 1$ model in four dimensions. Hence dimensional reduction and partial supersymmetry breaking are both present in this scenario. A technical tool that satisfies both features simultaneously is provided by the Scherk-Schwarz mechanism [1] (also called generalized dimensional reduction), where nontrivial boundary conditions for the fields along the compactified dimensions can lead to spontaneously broken gauge and space-time symmetries. In this paper the simplest setup to investigate partial supersymmetry breaking by generalized dimensional reduction is used, namely the partial breaking of $N = 2 \rightarrow N = 1$ in four dimensions coming from a theory in five dimensions. The focus is on the superHiggs effect, where Goldstone fermions from the broken supersymmetries become the longitudinal components of massive gravitinos.

The physics that underlies this superHiggs effect for the spontaneous breaking of $N = 2$ supersymmetry to $N = 1$ was investigated in a flat Minkowski background [2] as well as in an anti-de Sitter (AdS) background [3], where theories were constructed that describe partial supersymmetry breaking in a model-independent approach with a minimal field content motivated by the superHiggs effect. In this paper these theories of partial $N = 2 \rightarrow N = 1$ supersymmetry breaking in four dimensions will be reproduced by compactifying the massless $N = 2$ $D = 5$ massless gravitino multiplet¹ on the orbifold S^1/Z_2 and using the Scherk-Schwarz mechanism [1] to introduce a symmetry-breaking mass parameter.² In order to exhibit the superHiggs effect, this procedure must be carried out in a supergravity background.

This derivation serves several purposes: The starting point of the investigation is the massless $N = 2$ $D = 5$ gravitino multiplet Noether-coupled to $N = 2$ $D = 5$ supergravity which leads to pure $N = 4$ $D = 5$ supergravity (without matter). This is a rather simple theory that allows for easy extensions such as embedding into a higher N theory or introducing matter couplings. Second, the theories obtained from five dimensions are already out of unitary gauge, so that the symmetry breaking mass parameter m can be set to zero with impunity. In particular, no singular Stückelberg redefinitions (containing $1/m$ -terms as in Ref. [2]) are required.

What other motivation apart from simplicity is there to look at a theory in five dimensions? It seems natural to derive a massive 4-dimensional theory from a massless theory in five dimensions, because the degrees of freedom of a massless $D = 5$ vector field (3) and the degrees of freedom of a massless symplectic $D = 5$ Majorana spinor (4) match those of their massive counterparts in $D = 4$. This transition can be easily implemented by a Scherk-Schwarz compactification on S^1 , where it can be seen that the fifth compo-

¹ Here, $N = 2$ refers to the lowest supersymmetry in five dimensions, corresponding to two symplectic Majorana spinors, which are equivalent to one Dirac spinor.

²In Ref. [4] such a combination of orbifold projection and Scherk-Schwarz compactification was introduced to derive string theories with spontaneously broken supersymmetry.

nents of a symplectic gravitino $\Psi_{\alpha M}^i$ and a complex vector A_M , $\Psi_{\alpha 4}^i$ and A_4 , provide the longitudinal components of the massive 4-dimensional fields $\Psi_{\alpha m}^i$ and A_m .³

A natural candidate for partial supersymmetry breaking in four dimensions effected by Scherk-Schwarz compactification on S^1 would be pure $N = 2$ $D = 5$ supergravity [5, 6], in particular since the degrees of freedom exactly match those of one of the minimal cases (with two vectors) considered in Ref. [2]. Unfortunately, Scherk-Schwarz compactification applied to this theory breaks both supersymmetries. In fact, any Scherk-Schwarz compactification of N -extended supergravity on S^1 breaks an even number of supersymmetries, since the Scherk-Schwarz generator must be a generator of $USp(N)$ [7]. This can be circumvented by projecting out half of the states by compactifying on the orbifold S^1/Z_2 . In this paper, the length of the interval S^1/Z_2 is assumed to be of the order of the Planck length; hence only the massless Kaluza-Klein modes (possibly lifted by the Scherk-Schwarz mass parameter) are retained.

The outline of the paper is as follows: In the second section the massless $N = 2$ $D = 5$ gravitino multiplet will be compactified on S^1/Z_2 , yielding a massive $N = 1$ gravitino multiplet in four dimensions together with a spontaneously broken fermionic gauge symmetry. In the third section, the massless $N = 2$ $D = 5$ gravitino multiplet will be Noether-coupled to $N = 2$ $D = 5$ supergravity thus yielding $N = 4$ supergravity before the dimensional reduction is carried out. The corresponding 4-dimensional theory is shown to coincide with theories exhibiting partial supersymmetry breaking which were derived from four dimensions [8, 9, 10].⁴

2 Generalized compactification of the massless $N = 2$ $D = 5$ gravitino multiplet

The massless $N = 2$ $D = 5$ gravitino multiplet contains two symplectic Majorana gravitinos Ψ_{iM} , two symplectic Majorana spinors Λ_i and four real vector fields. Its Lagrangian and supersymmetry transformations can be conveniently obtained by truncation of the $N = 4$ $D = 5$ supergravity constructed in Ref. [14]. The conventions for the 5-dimensional Dirac algebra and the symplectic geometry are presented in the Appendix.

The five abelian vector fields A_M^{ij} $i, j \in \{1, \dots, 4\}$ in the 5-dimensional representation of $USp(4)$ which are part of the spectrum of $N = 4$ $D = 5$ supergravity satisfy the reality condition [5]

$$A_M^{ij} = (A_{Mij})^* ,$$

³In this paper, $i, j, \dots$ denote symplectic indices, $M, N, \dots$ are 5-dimensional world indices and $m, n, \dots$ are 4-dimensional world indices. A dotted numerical index stands for a world index, to be distinguished from undotted Lorentz indices.

⁴The geometric construction of [10] could be generalized to accommodate an arbitrary $N = 2$ matter and gauge content [11] and to give masses to light mirror fermions [12]. String models exhibiting the mass spectrum of partially broken supersymmetry were constructed in Refs. [13].

and are parametrized by

$$A_{Mij} = \begin{pmatrix} 0 & A_M & \bar{B}_M & \bar{C}_M \\ -A_M & 0 & -C_M & B_M \\ -\bar{B}_M & C_M & 0 & -A_M \\ -\bar{C}_M & -B_M & A_M & 0 \end{pmatrix}, \quad (1)$$

where A_M is real and B_M and C_M are complex.

Truncation of the $N = 4$ $D = 5$ supergravity to the massless $N = 2$ gravitino multiplet requires $A_M = 0$. The remaining two complex vectors can be written as

$$A_{Mia} = \begin{pmatrix} -\bar{B}_M & C_M \\ -\bar{C}_M & -B_M \end{pmatrix} \quad a, i \in \{1, 2\},$$

which corresponds to a decomposition $USp(4) \rightarrow USp(2) \otimes USp(2) \cong SU(2) \otimes SU(2)$. The field A_M now carries two different symplectic indices i and a , indicating the two independent $SU(2)$'s that it is charged under. Here,

$$\Omega_{ij} = \Omega_{ab} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

The Lagrangian is given by

$$\mathcal{L} = \frac{i}{2} \bar{\Psi}_{iM} \Gamma^{MNO} \partial_N \Psi_O^i - \frac{i}{2} \bar{\Lambda}_i \Gamma^M \partial_M \Lambda^i + \frac{1}{8} F_{MN}{}^i{}_a F^{MN}{}^a{}_i, \quad (2)$$

and the supersymmetry transformations are

$$\begin{aligned} \delta_{\Xi} \Psi_M^i &= \frac{1}{6} F_{NO}{}^i{}_a (\Gamma_M{}^{NO} + 2\delta_M^{[N} \Gamma^{O]}) \Xi^a \\ \delta_{\Xi} A_{Ma}{}^i &= -\frac{2i}{\sqrt{3}} \bar{\Xi}_a \Gamma_M \Lambda^i - 2i \bar{\Xi}_a \Psi_M^i \\ \delta_{\Xi} \Lambda^i &= -\frac{1}{2\sqrt{3}} F_{MN}{}^i{}_a \Gamma^{MN} \Xi^a, \end{aligned} \quad (3)$$

where $F_{MN}{}^i{}_a = \partial_{[M} A_{N]}^i{}_a$. In order to implement the Scherk-Schwarz mechanism on the orbifold S^1/Z_2 , the Z_2 transformation and the generator T of a global symmetry of the 5-dimensional theory used for the Scherk-Schwarz mechanism must satisfy [15]

$$Z_2 e^{imTx^4} = e^{imT(-x^4)} Z_2 \Leftrightarrow \{Z_2, T\} = 0. \quad (4)$$

In addition, the parity operation must be part of the discrete symmetries of the 5-dimensional theory. The effect of the parity operator $Z_2 = \pm \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \otimes i\Gamma^4$ on the fields is listed in Tables 1 and 2. With the choice $T = \sigma^2 \in su(2)$ and the transformation

$$\Phi^i \rightarrow (e^{im\sigma^2 x^4})^i{}_j \Phi^j$$

for a generic field Φ^i having a symplectic i (not a) index, the generalized dimensional reduction of the massless gravitino multiplet is completely specified.

	λ^i	ψ_m^i	ψ_4^i	ξ^a	i/a
Parity	1	-1	1	1	1
	-1	1	-1	-1	2

Table 1: Fermionic parity assignment of $D = 5$ $N = 2$ gravitino multiplet in terms of $D = 4$ Weyl spinors.

	B_m	B_4	C_m	C_4
Parity	-1	1	1	-1

Table 2: Bosonic parity assignment of $D = 5$ $N = 2$ gravitino multiplet in terms of $D = 4$ fields.

The limit $l \rightarrow 0$ of the length l of the interval S^1/Z_2 is implicitly understood, so that only the lowest even Fourier-modes of the fields which depend only on the first four space-time co-ordinates $x^0, \dots, x^3$ (zero-modes) need to be retained.

To simplify the identification of the physical fields in the 4-dimensional theory, the following redefinitions are necessary to diagonalize and normalize the fermionic kinetic terms:

$$\Psi_m^i \rightarrow \Psi_m^i + \frac{1}{2}\Gamma_m\Gamma^4\Psi_4^i, \quad (5)$$

and $\nu^i = \sqrt{3/2}\psi_4^i$. Note that in (5) no terms $\sim 1/m\partial_m\Psi_4^i$ as in Ref. [2] occur; in a gravitational background, the covariantized form of this term would not leave the gravitino kinetic term invariant and would therefore induce terms $\sim 1/m$ in the Lagrangian.

With additional “chiral” redefinitions $\Lambda^i \rightarrow -\Gamma^4\Lambda^i$ and $\Xi^a \rightarrow \Gamma^4\Xi^a$ the Lagrangian becomes

$$\begin{aligned} \mathcal{L}_4 = & \epsilon^{mnpq}\bar{\psi}_{2m}\bar{\sigma}_n\partial_p\psi_q^2 - i\bar{\lambda}_1\bar{\sigma}^m\partial_m\lambda^1 - i\bar{\nu}_1\bar{\sigma}^m\partial_m\nu^1 \\ & - \frac{1}{4}C_{mn}\bar{C}^{mn} - \frac{1}{2}\mathcal{D}_m B_4\mathcal{D}^m\bar{B}_4 \\ & - \sqrt{\frac{3}{2}}m(i\psi_m^2\sigma^m\bar{\nu}_1 + h.c.) - m(\nu^1\nu^1 + h.c.) \\ & + \frac{1}{2}m\lambda^1\lambda^1 - m\psi_m^2\sigma^{mn}\psi_n^2 + h.c. \end{aligned} \quad (6)$$

with supersymmetry transformations

$$\begin{aligned} \delta_\xi C_m &= i\frac{2}{\sqrt{3}}\xi^1\sigma_m\bar{\lambda}_1 + 2\xi^1\psi_m^2 - i\sqrt{\frac{2}{3}}\xi^1\sigma_m\bar{\nu}_1 \\ \delta_\xi B_4 &= \frac{2}{\sqrt{3}}\xi^1\lambda^1 + \frac{2\sqrt{2}}{\sqrt{3}}\xi^1\nu^1 \\ \delta_\xi\lambda^1 &= \frac{1}{\sqrt{3}}\bar{C}_{mn}\sigma^{mn}\xi^1 + \frac{i}{\sqrt{3}}\mathcal{D}_m B_4\sigma^m\bar{\xi}_1 \\ \delta_\xi\psi_m^2 &= -\frac{i}{2}C_{+mn}\sigma^n\bar{\xi}_1 + \mathcal{D}_m\bar{B}_4\xi^1 \\ \delta_\xi\nu^1 &= -\frac{1}{\sqrt{6}}\bar{C}_{mn}\sigma^{mn}\xi^1 + i\sqrt{\frac{2}{3}}\mathcal{D}_m B_4\sigma^m\bar{\xi}_1, \end{aligned}$$

where $\mathcal{D}_m B_4 = \partial_m B_4 - m C_m$ and $C_{+mn} = \partial_{[m} C_{n]} + \frac{i}{2} \epsilon_{mnrs} \partial^{[r} C^{s]}$. The expression for the gauge invariant derivative $\mathcal{D}_m$ shows that the complex scalar B_4 is the Goldstone boson of a spontaneously broken abelian gauge symmetry mediated by the vector C_m . The spinor ν^1 is the longitudinal component of the massive gravitino ψ_m^2 . The massless five-dimensional theory also contains a fermionic gauge symmetry:

$$\delta_\Theta \Psi_M^i = \frac{2}{\kappa} \partial_M \Theta^i .$$

A generalized Scherk-Schwarz compactification leads to a spontaneously broken fermionic symmetry:

$$\begin{aligned} \delta_\theta \psi_m^2 &= i \frac{m}{\kappa} \sigma_m \bar{\theta}_2 \\ \delta_\theta \nu^1 &= \sqrt{6} \frac{m}{\kappa} \theta^2 . \end{aligned} \quad (7)$$

Therefore, ν^1 can also be interpreted as the Goldstino for this broken symmetry.

The Lagrangian for the massless gravitino multiplet (2) is not unique. In flat 5-dimensional space-time a massless vector B_M is dual⁵ to an antisymmetric tensor G_{MN}

$$\partial_{[M} B_{N]} = \frac{1}{2} \epsilon_{MNOPQ} \partial^O G^{PQ} + \dots, \quad (8)$$

where the dots stand for contributions from interaction terms. So all four real vectors of the massless $N = 2$ gravitino multiplet can be dualized and then dimensionally reduced to four dimensions. The implementation of the Scherk-Schwarz mechanism on S^1/Z_2 requires that fields of the same index structure come in pairs, so that one of them can be projected out by the Z_2 reflection whereas the other becomes a massive field in the 4-dimensional theory. If both B_M and C_M are dualized, the field strength $F_{MN}^i{}_a$ is simply replaced by its dual in the Lagrangian and the supersymmetry transformations and the automorphism group decomposes as before as $USp(4) \rightarrow SU(2) \otimes SU(2)$.

The other possibility is the dualization of only two vectors, which must be the real or imaginary parts of the vectors $B_M = B_M^R + i B_M^I$ and $C_M = C_M^R + i C_M^I$. Otherwise, either the complex vector or the complex antisymmetric tensor are completely projected out. Here, the imaginary parts are chosen to be dualized: $B_M^I \rightarrow B_{MN}^I$ and $C_M^I \rightarrow C_{MN}^I$. This corresponds to the decomposition

$$\begin{aligned} F_{MN}^i{}_a &= \text{Re}(F_{MN}^i{}_a) + i \text{Im}(F_{MN}^i{}_a) \\ &\rightarrow F_{MN}^R{}^i{}_a + i v_{MN}^i{}_a , \end{aligned}$$

where $v_{MN}^i{}_a = 1/2 \epsilon_{MNOPQ} \partial^O G^{PQ}{}^i{}_a$ and $G_{MN}^i{}_a = \begin{pmatrix} -C_{MN}^I & B_{MN}^I \\ B_{MN}^I & C_{MN}^I \end{pmatrix}$. To conserve this structure, the $SU(2) \otimes SU(2)$ transformations must be restricted to those which do not mix real and imaginary parts. Therefore only the generator σ^2 is allowed, which corresponds to a decomposition $SU(2) \otimes SU(2) \rightarrow U(1) \otimes U(1)$.

⁵This is not true for a 5-dimensional AdS background, where antisymmetric tensors satisfy additional self-duality conditions [16].

	B_m^R	B_4^R	C_m^R	C_4^R	B_{mn}^I	B_{m4}^I	C_{mn}^I	C_{m4}^I
Parity	-1	1	1	-1	1	-1	-1	1

Table 3: Bosonic parity assignment of the dualized $D = 5$ $N = 2$ gravitino multiplet in terms of $D = 4$ fields.

Dualization of the imaginary parts of the complex vectors B_M and C_M yields the Lagrangian

$$\begin{aligned} \mathcal{L} = & \frac{i}{2} \bar{\Psi}_{iM} \Gamma^{MNO} \partial_N \Psi_O^i - \frac{i}{2} \bar{\chi}_i \Gamma^M \partial_M \chi^i \\ & + \frac{1}{8} F_{MN}^R{}^i F^{RMNa}{}_i + \frac{1}{8} v_{MN}^i v^{MNa}{}_i, \end{aligned} \quad (9)$$

and the supersymmetry transformations are

$$\begin{aligned} \delta_{\Xi} \Psi_M^i &= \frac{1}{6} F_{NO}^R{}^i (\Gamma_M^{NO} + 2\delta_M^{[N} \Gamma^{O]}) \Xi^a + \frac{i}{6} v_{NO}^i (\Gamma_M^{NO} + 2\delta_M^{[N} \Gamma^{O]}) \Xi^a \\ \delta_{\Xi} A_{Ma}^R{}^i &= -\frac{i}{\sqrt{3}} \bar{\Xi}_a \Gamma_M \Lambda^i - i \bar{\Xi}_a \Psi_M^i + h.c. \\ \delta_{\Xi} G_{MNa}^i &= \frac{1}{\sqrt{3}} \bar{\Xi}_a \Gamma_{MN} \chi^i - \bar{\Xi}_a \Gamma_{[M} \Psi_{N]}^i + h.c. \\ \delta_{\Xi} \Lambda^i &= -\frac{1}{2\sqrt{3}} F_{MN}^R{}^i \Gamma^{MN} \Xi^a - \frac{i}{2\sqrt{3}} v_{MN}^i \Gamma^{MN} \Xi^a. \end{aligned} \quad (10)$$

The parities of the antisymmetric tensors in terms of 4-dimensional fields are determined by the parities of the dual vectors and the dualization relation (8) and are listed in Table 3. Performing the same procedure and redefinitions (5) as before, the following Lagrangian is obtained:

$$\begin{aligned} \mathcal{L}_4 = & \epsilon^{mnpq} \bar{\psi}_{2m} \bar{\sigma}_n \partial_p \psi_q^2 - i \bar{\lambda}_1 \bar{\sigma}^m \partial_m \lambda^1 - i \bar{\nu}_1 \bar{\sigma}^m \partial_m \nu^1 \\ & - \frac{1}{4} C_{mn}^R C^{Rmn} - \frac{1}{4} \mathcal{F}_{mn}^{C_4^I} \mathcal{F}^{C_4^I mn} + \frac{1}{4} v_m^{B^I} v^{B^I m} - \frac{1}{2} \mathcal{D}_m B_4^R \mathcal{D}^m B_4^R \\ & - \sqrt{\frac{3}{2}} m (i \psi_m^2 \sigma^m \bar{\nu}_1 + h.c.) - m (\nu^1 \nu^1 + h.c.) \\ & + \frac{1}{2} m \lambda^1 \lambda^1 - m \psi_m^2 \sigma^{mn} \psi_n^2 + h.c. \end{aligned} \quad (11)$$

with supersymmetry transformations

$$\begin{aligned} \delta_{\xi} C_m^R &= i \frac{1}{\sqrt{3}} \xi^1 \sigma_m \bar{\lambda}_1 + \xi^1 \psi_m^2 - i \frac{1}{\sqrt{6}} \xi^1 \sigma_m \bar{\nu}_1 + h.c. \\ \delta_{\xi} B_4^R &= \frac{1}{\sqrt{3}} \xi^1 \lambda^1 + \sqrt{\frac{2}{3}} \xi^1 \nu^1 + h.c. \\ \delta_{\xi} B_{mn}^I &= -\frac{2}{\sqrt{3}} \xi^1 \sigma_{mn} \lambda^1 - 2 \sqrt{\frac{2}{3}} \xi^1 \sigma_{mn} \nu^1 - i \xi^1 \sigma_{[m} \bar{\psi}_{2n]} + h.c. \end{aligned}$$

$$\begin{aligned}
\delta_\xi C_{m4}^I &= -\frac{i}{\sqrt{3}}\xi^1\sigma_m\bar{\lambda}_1 + \frac{i}{\sqrt{6}}\xi^1\sigma_m\bar{\nu}_1 + \xi^1\psi_m^2 + h.c. \\
\delta_\xi\lambda^1 &= \frac{1}{\sqrt{3}}C_{mn}^R\sigma^{mn}\xi^1 + \frac{i}{\sqrt{3}}\mathcal{D}_m B_4^R\sigma^m\bar{\xi}_1 - \frac{1}{\sqrt{3}}\mathcal{F}_{mn}^{C_4^I}\sigma^{mn}\xi^1 + \frac{1}{\sqrt{3}}v_m^{B^I}\sigma^m\bar{\xi}_1 \\
\delta_\xi\nu^1 &= -\frac{1}{\sqrt{6}}C_{mn}^R\sigma^{mn}\xi^1 + i\sqrt{\frac{2}{3}}\mathcal{D}_m B_4^R\sigma^m\bar{\xi}_1 + \frac{1}{\sqrt{6}}\mathcal{F}_{mn}^{C_4^I}\sigma^{mn}\xi^1 + \sqrt{\frac{2}{3}}v_m^{B^I}\sigma^m\bar{\xi}_1 \\
\delta_\xi\psi_m^2 &= -\frac{i}{2}C_{+mn}^R\sigma^n\bar{\xi}_1 + \mathcal{D}_m B_4^R\xi^1 - \frac{i}{2}\mathcal{F}_{+mn}^{C_4^I}\sigma^n\bar{\xi}_1 + iv_m^{B^I}\xi^1,
\end{aligned}$$

where $\mathcal{D}_m B_4^R = \partial_m B_4^R - mC_m^R$, $\mathcal{F}_{mn}^{C_4^I} = \partial_{[m}C_{n]4}^I - mB_{mn}^I$ and $v_m^{B^I} = 1/2\epsilon_{mnop}\partial^n B^{Iop}$. In the dual case, the vector C_{m4}^I becomes the Goldstone boson of the spontaneously broken gauge symmetry mediated by the antisymmetric tensor B_{mn}^I . This is the dual Higgs mechanism investigated in Ref. [17]. As before, the 5-dimensional theory has a fermionic gauge symmetry which is spontaneously broken upon generalized dimensional reduction (7). The generalized dimensional reduction of the massless gravitino multiplet with four antisymmetric tensors is analogous to the case illustrated above and will not be presented here.

These 4-dimensional theories correspond exactly, up to field relabelings, to the dual versions of the massive $N = 1$ gravitino multiplet (out of unitary gauge) before being coupled to gravity [2]. In order to exhibit the superHiggs effect for partial supersymmetry breaking, the above multiplets must be coupled to gravity and the fermionic gauge symmetry must be promoted to a local supersymmetry. This will be addressed in the next section.

3 Generalized dimensional reduction of pure $N = 4$ $D = 5$ supergravity

The complete theory of partially broken supergravity in four dimensions to all orders in the fields should now be obtainable by Noether-coupling the massless gravitino multiplet to pure $N = 2$ $D = 5$ supergravity [5, 6, 18] and performing a generalized dimensional reduction. Since consistency requires that the fermionic gauge symmetries of the massless gravitinos become local supersymmetries, the resulting theory must be $N = 4$ supersymmetric. Therefore, the Scherk-Schwarz mechanism should be applied to $N = 4$ $D = 5$ supergravity. The pure five-dimensional $N = 4$ supergravity was shown to be derivable as a consistent truncation of the maximal $N = 8$ supergravity [5]. The explicit form of the $N = 4$ supergravity Lagrangian and its supersymmetry transformations was given in Ref. [14].

For the sake of clarity representative parts of the result of [14] are reproduced⁶. $N = 4$ supergravity has a $USp(4)$ symmetry inherited from the automorphism group of the supersymmetry algebra and its field content together with a consistent parity assignment

⁶Here, the vector in the singlet representation of $USp(4)$, B_μ , from Ref. [14] has been renamed to G_M and the vector fields in the 5-dimensional representation of $USp(4)$ are parametrized as in (1). Also, the 5-dimensional spinors χ^i have been renamed to Λ^i . The following redefinitions were performed: $\Gamma^A \rightarrow -i\Gamma^A$, $A_{Mij} \rightarrow 1/2A_{Mij}$, and $\Xi^i \rightarrow 2\Xi^i$.

	λ^i	ψ_m^i	ψ_4^i	ξ^i	i
Parity	-1	1	-1	1	1
	1	-1	1	-1	2
	1	-1	1	-1	3
	-1	1	-1	1	4

Table 4: Fermionic fields and parities of $D = 5$ $N = 4$ supergravity in terms of $D = 4$ Weyl spinors.

	e_m^a	e_m^4	e_4^4	G_m	G_4	σ	A_m	A_4	B_m	B_4	C_m	C_4
Parity	1	-1	1	-1	1	1	-1	1	-1	1	1	-1
	$N = 2$ supergravity					vector mult.			gravitino mult.			

Table 5: Bosonic fields and parities of $D = 5$ $N = 4$ supergravity in terms of $D = 4$ fields.

for inversion of the fifth co-ordinate is given in Tables 4 and 5. The Lagrangian is given up to terms of order $\mathcal{O}(\kappa)\mathcal{O}(\text{fermions})^7$ and the supersymmetry transformations are given up to three-fermion terms. The Lagrangian reads

$$\begin{aligned}
\kappa e^{-1}\mathcal{L} = & -\frac{1}{2\kappa^2}\mathcal{R} + \frac{i}{2}\bar{\Psi}_{iM}\Gamma^{MNO}D_N\Psi_O^i - \frac{i}{2}\bar{\Lambda}_i\Gamma^M D_M\Lambda^i \\
& -\frac{1}{16}e^{\frac{2}{\sqrt{3}}\kappa\sigma}F_{MN}^{ij}F_{ij}^{MN} - \frac{1}{4}e^{-\frac{4}{\sqrt{3}}\kappa\sigma}G_{MN}G^{MN} - \frac{1}{2}\partial_M\sigma\partial^M\sigma \\
& + \frac{\kappa}{16\sqrt{2}}e^{-1}\epsilon^{MNOPQ}F_{MN}^{ij}F_{OPij}G_Q + \mathcal{O}(\kappa)\mathcal{O}(\text{fermions}) , \tag{12}
\end{aligned}$$

and the supersymmetry transformations are

$$\begin{aligned}
\delta_{\Xi}\Psi_{Mi} &= \frac{2}{\kappa}D_M\Xi_i - \frac{1}{6}e^{\frac{1}{\sqrt{3}}\kappa\sigma}F_{NOij}(\Gamma_M^{NO} + 2\delta_M^{[N}\Gamma^{O]})\Xi^j \\
&\quad - \frac{1}{6\sqrt{2}}e^{-\frac{2}{\sqrt{3}}\kappa\sigma}G_{NO}(\Gamma_M^{NO} + 2\delta_M^{[N}\Gamma^{O]})\Xi_i + (3\text{-fermion terms}) \\
\delta_{\Xi}A_M^{ij} &= \frac{i}{\sqrt{3}}e^{-\frac{1}{\sqrt{3}}\kappa\sigma}(2\bar{\Xi}^{[i}\Gamma_M\Lambda^{j]} - \Omega^{ij}\bar{\Xi}_k\Gamma_M\Lambda^k) \\
&\quad - ie^{-\frac{1}{\sqrt{3}}\kappa\sigma}(2\bar{\Xi}^{[i}\Psi_M^{j]} - \Omega^{ij}\bar{\Xi}_k\Psi_M^k) \\
\delta_{\Xi}\Lambda_i &= -\Gamma^M\Xi_i\partial_M\sigma - \frac{1}{2\sqrt{3}}e^{\frac{1}{\sqrt{3}}\kappa\sigma}F_{MNij}\Gamma^{MN}\Xi^j \\
&\quad + \frac{1}{\sqrt{6}}e^{\frac{2}{\sqrt{3}}\kappa\sigma}G_{MN}\Gamma^{MN}\Xi_i + (3\text{-fermion terms}) \\
\delta_{\Xi}G_M &= i\sqrt{\frac{2}{3}}e^{\frac{2}{\sqrt{3}}\kappa\sigma}\bar{\Xi}_i\Gamma_M\Lambda^i + \frac{i}{\sqrt{2}}e^{\frac{2}{\sqrt{3}}\kappa\sigma}\bar{\Xi}_i\Psi_M^i \\
\delta_{\Xi}\sigma &= -i\bar{\Xi}_i\Lambda^i \\
\delta_{\Xi}e^A_M &= i\kappa\bar{\Xi}_i\Gamma^A\Psi_M^i . \tag{13}
\end{aligned}$$

⁷The Lagrangian derived in Ref. [14] includes such terms up to four-fermion terms, but they are not illustrative in the context of partial supersymmetry breaking.

The fields and supersymmetry parameters that are enclosed in double lines in Tables 4 and 5 are the ones coming from the gravitino multiplet. This field content shows that a consistent Noether-coupling of the gravitino multiplet to $N = 2$ supergravity necessitates the inclusion of a vector multiplet with fields $(\lambda^1, \lambda^2, \sigma, A_m, A_4)$.

From the parities in Tables 4 and 5 it is clear that the four dimensional theory contains in addition to the fields of $N = 2$ $D = 4$ supergravity and the massive $N = 1$ gravitino multiplet out of unitary gauge [2] two chiral multiplets $(\lambda^2, \psi_4^2, P, G_4, \sigma, A_4)$ where $e^P = e^4_4$. This is the field content of one vector- and one hyper-multiplet coupled to $N = 2$ supergravity.

The rest of this section is now devoted to showing that a generalized dimensional reduction of $N = 4$ $D = 5$ supergravity on S^1/Z_2 with appropriately chosen $USp(4)$ -generator for the Scherk-Schwarz mechanism indeed leads to partially broken $N = 2$ supergravity in four dimensions. First, those $USp(4)$ -generators which satisfy the condition (4) with $Z_2 = \pm \text{diag}(1, -1, -1, 1) \otimes i\Gamma^4$ must be singled out. Out of T_r $r \in \{1, \dots, 10\}$ (see Appendix), the generators T_s with $s \in \{1, 2, 4, 5, 8, 10\}$ satisfy (4).

From the supersymmetry transformation of the gravitinos (13)

$$\begin{aligned} \delta_\Xi \Psi_{iM} &= \frac{2}{\kappa} D_M \Xi_i + \dots \\ \rightarrow \quad \delta_\Xi (e^{imT_s x^4})^j_i \Psi_{iM} &= \frac{2}{\kappa} D_M \left((e^{imT_s x^4})^j_i \Xi_j \right) + \dots \end{aligned} \quad (14)$$

it is obvious that the fifth component Ψ_{i4} of the gravitinos can pick up a constant shift $\sim m$. With the redefinition (5) the corresponding 4-dimensional gravitino Ψ_{im} will also pick up a shift, thus identifying the broken supersymmetries.

The choice T_s with $s \in \{8, 10\}$ would break all supersymmetries. In order to match the Scherk-Schwarz mechanism in Sec. 2, the generator $T_5 = \begin{pmatrix} 0 & 0 \\ 0 & \sigma^2 \end{pmatrix}$ will be used to implement the partial breaking of supersymmetry in the 4-dimensional theory. With T_5 , both ψ_4^3 and ψ_4^4 acquire a shift under supersymmetry transformations

$$\delta_\xi \begin{pmatrix} \psi_4^3 \\ \psi_4^4 \end{pmatrix} = \frac{2}{\kappa} m \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} \xi^3 \\ \xi^4 \end{pmatrix} + \dots$$

Since ψ_4^4 is projected out under the Z_2 reflection, only ψ_4^3 will be the Goldstino of one spontaneously broken supersymmetry in the 4-dimensional theory.

The dimensional reduction from $M_5 \rightarrow M_4 \times S^1/Z_2$ requires Weyl rescalings and field redefinitions described in Refs. [6, 19] in order to obtain canonically normalized diagonal kinetic terms. They are simplified, however, because the vector e^4_m from the fünfbein used for the Weyl rescaling is odd under the Z_2 reflection. Explicitly the rescalings and redefinitions are: $e^a_m \rightarrow e^{-P/2} e^a_m$, $\xi^i \rightarrow e^{-P/4} \xi^i$, $\lambda^i \rightarrow -i e^{P/4} \lambda^i$, $\psi_m^1 \rightarrow e^{-P/4} \psi_m^1 + i/2 \sigma_m \bar{\psi}_{2,4} e^{-3P/2}$, $\psi_m^4 \rightarrow e^{-P/4} \psi_m^4 - i/2 \sigma_m \bar{\psi}_{3,4} e^{-3P/2}$, $\psi_4^i \rightarrow e^{5P/4} \sqrt{2/3} \nu^i$.

The resulting 4-dimensional Lagrangian up to terms of order $\mathcal{O}(\kappa)\mathcal{O}(\text{fermions})$ in the conventions of [20] reads

$$\begin{aligned} e^{-1} \mathcal{L} = & \\ & - \frac{1}{2\kappa^2} \mathcal{R} + \epsilon^{pqrs} \bar{\psi}_{ip} \bar{\sigma}_q D_r \psi_s^i - i \bar{\chi}_i \bar{\sigma}^m D_m \chi^i - i \bar{\Omega}_i \bar{\sigma}^m D_m \Omega^i \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{4}e^{\sqrt{2}\kappa\varphi}\mathcal{A}_{mn}\bar{\mathcal{A}}^{mn}-\frac{1}{2}\partial_m\varphi\partial^m\varphi-\frac{1}{2}e^{-2\sqrt{2}\kappa\varphi}\partial_m\pi\partial^m\pi-\frac{1}{2}\partial_m\tilde{\varphi}\partial^m\tilde{\varphi}-\frac{1}{2}e^{2\kappa\tilde{\varphi}}\mathcal{D}_m\pi^a\mathcal{D}^m\pi^a \\
& -me^{\kappa(\tilde{\varphi}-\frac{1}{\sqrt{2}}\varphi)}\left(-\frac{i}{\sqrt{2}}\psi_m^2\sigma^m\bar{\Omega}_1+i\psi_m^2\sigma^m\bar{\chi}_1-\sqrt{2}\Omega^1\chi^1+\frac{1}{2}\chi^1\chi^1+\psi_m^2\sigma^{mn}\psi_n^2+h.c.\right) \\
& +\frac{\kappa}{4\sqrt{2}}\pi\tilde{\mathcal{A}}_{mn}\bar{\mathcal{A}}^{mn}+\mathcal{O}(\kappa)\mathcal{O}(\text{fermions})\,,
\end{aligned} \tag{15}$$

and the supersymmetry transformations up to three-fermion terms are

$$\begin{aligned}
\delta_\xi e^a_m &= -i\kappa\psi_m^i\sigma^a\bar{\xi}_i+h.c. \\
\delta_\xi\psi_m^i &= \frac{2}{\kappa}D_m\xi^i-\frac{1}{2}\epsilon^{ij}e^{\kappa\frac{\varphi}{\sqrt{2}}}\mathcal{A}_{+mn}\sigma^n\bar{\xi}_j+\frac{i}{\sqrt{2}}e^{-\sqrt{2}\kappa\varphi}\partial_m\pi\xi^i \\
& -ie^{\kappa\tilde{\varphi}}\mathcal{D}_m\pi^a\sigma^{ai}_j\xi^j+i\frac{m}{\kappa}\delta_2^ie^{\kappa(\tilde{\varphi}-\frac{1}{\sqrt{2}}\varphi)}\sigma_m\bar{\xi}_2 \\
& +(3\text{-fermion terms}) \\
\delta_\xi\mathcal{A}_m &= e^{-\kappa\frac{\varphi}{\sqrt{2}}}(-2i\epsilon_{ij}\psi_m^i\xi^j-\sqrt{2}\bar{\Omega}_i\bar{\sigma}_m\xi^i) \\
\delta_\xi\Omega^i &= -\frac{i}{\sqrt{2}}e^{\kappa\frac{\varphi}{\sqrt{2}}}\bar{\mathcal{A}}_{mn}\sigma^{mn}\xi^i-i\epsilon^{ij}\sigma^m\bar{\xi}_j(\partial_m\varphi-ie^{-\sqrt{2}\kappa\varphi}\partial_m\pi) \\
& -\sqrt{2}\frac{m}{\kappa}\epsilon^{i2}e^{\kappa(\tilde{\varphi}-\frac{1}{\sqrt{2}}\varphi)}\xi^2+(3\text{-fermion terms}) \\
\delta_\xi\chi^i &= -i\epsilon^{ij}\sigma^m\bar{\eta}_j\partial_m\tilde{\varphi}-e^{\kappa\tilde{\varphi}}\epsilon^{ij}\sigma^a_j{}^k\mathcal{D}_m\pi^a\sigma^m\bar{\eta}_k \\
& +2\frac{m}{\kappa}\epsilon^{i2}e^{\kappa(\tilde{\varphi}-\frac{1}{\sqrt{2}}\varphi)}\xi^2+(3\text{-fermion terms}) \\
\delta_\xi\pi &= e^{\sqrt{2}\kappa\varphi}(i\epsilon_{ij}\Omega^i\xi^j+h.c.) \\
\delta_\xi\varphi &= \epsilon_{ij}\Omega^i\xi^j+h.c. \\
\delta_\xi\tilde{\varphi} &= \epsilon_{ij}\chi^i\xi^j+h.c. \\
\delta_\xi\pi^a &= e^{-\kappa\tilde{\varphi}}(-i\chi^i\sigma^a_i{}^j\epsilon_{jk}\xi^k+h.c.)\,,
\end{aligned} \tag{16}$$

where $\mathcal{D}_m\pi^a\sigma^a_i{}^j=\partial_m\pi^a\sigma^a_i{}^j-m(\sigma^1_i{}^jA_m+\sigma^2_i{}^jB_m)$, $\sigma^{ai}_j=\epsilon^{il}\epsilon_{jk}\sigma^a_l{}^k$, $\mathcal{A}_m=A_m+iB_m$, and $\tilde{\mathcal{A}}_{mn}=\epsilon_{mnop}\mathcal{A}^{op}$. Here, several field redefinitions have been performed to facilitate the identification of 4-dimensional $N=2$ multiplets. With the rescaling $P\rightarrow\sqrt{2/3}P$ the chart of fields in Eqs. (15, 16) corresponding to the fields in Eqs. (12, 13) is given in Table 6. In terms of $N=2$ multiplets, the fields $(e^a_m, \psi_m^1, \psi_m^2, A_m)$ constitute the $N=2$ supergravity multiplet, $(\varphi, \pi, \Omega^1, \Omega^2, B_m)$ constitute an $N=2$ vector multiplet, and $(\tilde{\varphi}, \pi^1, \pi^2, \pi^3, \chi^1, \chi^2)$ form a hyper-multiplet.

From the Lagrangian (15) and the supersymmetry transformations (16) it is obvious that ψ_m^2 is the gauge field for the spontaneously broken supersymmetry, whereas a linear combination of Ω^1 and χ^1 is the associated Goldstino. Its supersymmetric partner, the scalar $\pi^1+i\pi^2$ is the Goldstone boson of a spontaneously broken central charge gauged by the vector $\mathcal{A}_m$. That can be seen from the closure of the first and second supersymmetry algebra on $\pi^1+i\pi^2$ and $\mathcal{A}_m$ [2].

This result should be compared to partially broken 4-dimensional $N=2$ supergravities with complete $N=2$ multiplets constructed in Refs. [8, 9, 10]. Albeit equivalent up to field redefinitions, the non-singular parametrization of the scalar fields in Ref. [9] is closest to the parametrization of the scalars as obtained by the above-described dimensional

part. broken $N = 2$ $D = 4$ SUGRA	$N = 4$ $D = 5$ SUGRA
φ	$(\sqrt{2}\sigma + P)/\sqrt{3}$
$\tilde{\varphi}$	$(\sigma - \sqrt{2}P)/\sqrt{3}$
π	G_4
$\pi^1 + i\pi^2$	B_4
$-\pi^3$	A_4
$\mathcal{A}_m$	C_m
$e^a{}_m$	$e^a{}_m$
ψ_m^1	ψ_m^1
ψ_m^2	ψ_m^4
χ^{4-i}	$(\lambda^i + \sqrt{2}\nu^i)/\sqrt{3}$
Ω^{4-i}	$(\sqrt{2}\lambda^i - \nu^i)/\sqrt{3}$

Table 6: Corresponding fields of partially broken $N = 2$ $D = 4$ supergravity and pure $N = 4$ $D = 5$ supergravity.

reduction, so the field labels from [9] are used in Table 6. In Ref. [9] the breaking of the i th supersymmetry is parametrized by the quantities μ_i , $i \in \{1, 2\}$. In this context, the second supersymmetry is chosen to be broken, so $\mu_1 = 0$ and $\mu_2 = m$. The more general supersymmetry breaking scenario in Ref. [9] can be obtained from five dimensions by the choice $\mu_1 T_2 + \mu_2 T_5$ for the Scherk-Schwarz generator. One finds complete agreement of (15) and (16) up to trivial phase factors with the corresponding expressions in Ref. [9].

4 Conclusions

In this paper it was shown that partial supersymmetry breaking $N = 2 \rightarrow N = 1$ in four dimensions can be easily reproduced by compactifying $N = 4$ $D = 5$ supergravity on the orbifold S^1/Z_2 and using the Scherk-Schwarz mechanism. This means that compactification of $N = 4$ $D = 5$ supergravity on S^1/Z_2 automatically leads to an $N = 2$ supersymmetric theory in four dimensions in which no prepotential exists for the vector multiplet — thus allowing for partial supersymmetry breaking [10]. Although the derivation of partially broken theories in four dimensions from five dimensions is considered here only as a convenient tool, it allows for straightforward extensions like matter couplings or embeddings in higher- N theories in five dimensions. This is possible because the starting point — $N = 4$ $D = 5$ supergravity — is a linearly realized massless theory with complete multiplets. Therefore it is not surprising that complete $N = 2$ supermultiplets as in Refs. [8, 9, 10] persist in four dimensions in the nonlinearly realized broken phase.

The Poincaré dualities exploited in Ref. [2] for massless vectors and scalars in four dimensions can now be seen to be consequences of a duality in five dimension relating a massless vector to a massless antisymmetric tensor. Dual formulations of the theories in Refs. [8, 9, 10] can be found by dualizing first the Goldstone scalars to antisymmetric tensors and then dualizing the vectors according to the method described in Ref. [21]. These theories correspond to lowest order in the fields to the effective theories derived in Ref. [2].

In Ref. [3] the partial breaking of extended supersymmetry in four dimensions with a minimal field content as dictated by the superHiggs effect was extended to an anti-de Sitter background. There it was found that only one of the two Goldstone scalars in the theory out of unitary gauge could be dualized to an antisymmetric tensor, the other one could not be dualized because it occurred in the theory without derivatives acting upon it. If one were to derive this theory by a compactification $AdS_5 \rightarrow AdS_4 \times S^1/Z_2$, the resulting 4-dimensional theory would as well contain one vector and one antisymmetric tensor, since the bosonic group structure of the $N = 4$ AdS_5 automorphism group requires two of the five vectors, which reside in the 5-dimensional representation of $USp(4)$ in a Minkowski background, to be replaced by antisymmetric tensors [22]. The gauge group then becomes $SU(2) \otimes U(1)$ and the Z_2 reflection would project out two of the vectors in $SU(2)$ and one antisymmetric tensor in $U(1)$. Work along these lines is in progress.

Acknowledgments

I would like to thank the FNAL theory group for their hospitality during my stay as a summer visitor, when this work was initiated. I am also grateful to J. Bagger and Yi-Y. Wu for helpful discussions.

Appendix

The 5-dimensional Dirac algebra with the space-time metric $\eta_{AB} = \text{diag}(-1, 1, 1, 1, 1)$ reads

$$\{\Gamma^A, \Gamma^B\} = -2\eta^{AB} ,$$

where $\Gamma^A \in \{\gamma^0, \gamma^1, \gamma^2, \gamma^3, \gamma^5\}$. The matrices γ^a and γ^5 are defined as in Ref. [20] and the antisymmetric combinations $\Gamma^{A_1 \dots A_n} = \frac{1}{n!} \Gamma^{[A_1} \dots \Gamma^{A_n]}$ are defined with strength one. The 5-dimensional epsilon-tensor is defined by $\epsilon^{01234} = 1$.

The real symplectic metric to lower $USp(2N)$ -indices is chosen to be $\Omega_{ij} = 1_N \otimes i\sigma^2$. It is used to raise and lower symplectic indices of the vectors V^i and V_i according to $V^i = \Omega^{ij} V_j$ and $V_i = \Omega_{ij} V^j$.

A symplectic Majorana spinor in five dimensions (denoted by upper case Greek letters) is written in terms of Weyl spinors in four dimensions (denoted by lower case Greek letters) as

$$\Psi^i = \begin{pmatrix} \psi_\alpha^i \\ -\Omega^{ij} \bar{\psi}_j^{\dot{\alpha}} \end{pmatrix} , \quad \bar{\Psi}_i = (\Omega_{ij} \psi^{j\alpha} \quad \bar{\psi}_{i\dot{\alpha}}) ,$$

where $\bar{\psi}_i^{\dot{\alpha}} = \epsilon^{\dot{\alpha}\beta} (\psi_\beta^i)^*$. ‘‘Symplectic’’ indices on Weyl spinors in four dimensions are merely labels — they are not covariant indices.

The massless superalgebra in five dimensions has a $USp(N)$ automorphism group. $USp(N)$ with N even is the compact Lie group defined by the set of complex $N \times N$ matrices that are both unitary and symplectic. The generators of the corresponding Lie algebra $usp(N)$ form a set of $N(N+1)/2$ hermitian matrices T_r that satisfy the symplectic condition $T_r \Omega_{..} + \Omega_{..} T_r^\top = 0 \quad \forall r \in \{1, \dots, N(N+1)/2\}$. The representation of the basis

elements of $usp(4)$ used in this paper is

$$T_r \in \left\{ \begin{pmatrix} \sigma^j & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & \sigma^j \end{pmatrix}, \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -\sigma^2 \\ -\sigma^2 & 0 \end{pmatrix}, \begin{pmatrix} 0 & \sigma^3 \\ \sigma^3 & 0 \end{pmatrix}, \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} \right\}$$

with $r \in \{1, \dots, 10\}$.

References

- [1] J. Scherk and J. H. Schwarz, Phys. Lett. **B82** (1979) 60; Nucl. Phys. **B153** (1979) 61.
- [2] R. Altendorfer and J. Bagger, Phys. Lett. **B460** (1999) 127.
- [3] R. Altendorfer and J. Bagger, “*Dual Anti-de Sitter Superalgebras from Partial Supersymmetry Breaking*”, hep-th/9908084, to appear in Phys. Rev. **D**.
- [4] C. Kounnas and M. Porrati, Nucl. Phys. **B310** (1988) 355.
- [5] E. Cremmer, “*Supergravities in 5 dimensions*” in “*Superspace & Supergravity*”, ed. by S. W. Hawking and M. Roček, Cambridge University Press, 1981.
- [6] A. H. Chamseddine and H. Nicolai, Phys. Lett. **B96** (1980) 89.
- [7] E. Cremmer, J. Scherk, and J. H. Schwarz, Phys. Lett. **B84** (1979) 83.
- [8] S. Cecotti, L. Girardello, and M. Porrati, Phys. Lett. **B168** (1986) 83.
- [9] Yu. M. Zinov’ev, Sov. J. Nucl. Phys. **46** (1987) 540.
- [10] S. Ferrara, L. Girardello, and M. Porrati, Phys. Lett. **B366** (1996) 155.
- [11] P. Fré, L. Girardello, I. Pesando, and M. Trigiante, Nucl. Phys. **B493** (1997) 231.
- [12] L. Girardello, M. Porrati, and A. Zaffaroni, Nucl. Phys. **B505** (1997) 272.
- [13] E. Kiritsis and C. Kounnas, Nucl. Phys. **B503** (1997) 117;
I. Antoniadis, G. D’Appollonio, E. Dudas, and A. Sagnotti, Nucl. Phys. **B553** (1999) 133.
- [14] M. Awada and P. K. Townsend, Nucl. Phys. **B255** (1985) 617.
- [15] E. Dudas and C. Grojean, Nucl. Phys. **B507** (1997) 553.
- [16] M. Günaydin and N. Marcus, Class. Quant. Grav. **2** (1985) L11.
- [17] F. Quevedo and C. Trugenberger, Nucl. Phys. **B501** (1997) 143.
- [18] R. D’Auria, E. Maina, T. Regge, and P. Fré, Annals Phys. **135** (1981) 237.
- [19] E. Cremmer and B. Julia, Nucl. Phys. **B159** (1979) 141.

- [20] J. Wess and J. Bagger, “*Supersymmetry and Supergravity*”, Princeton University Press, 1992.
- [21] H. Nicolai and P. K. Townsend, Phys. Lett. **B98** (1981) 257.
- [22] L. J. Romans, Nucl. Phys. **B267** (1986) 433;
M. Günaydin, L. J. Romans, and N. P. Warner, Nucl. Phys. **B272** (1986) 598.